

A Review of Brain Tumor, Symptoms, Risk factors and Brain tumor in Children and Young Adults

Rehana*

*Institute of Molecular Biology and Biotechnology, Bahauddin Zakariya University, Multan-60800, Pakistan.

Article Received: 09 October 2019

Article Accepted: 17 December 2019

Article Published: 19 January 2020

ABSTRACT

A brain tumor rises when irregular cells form the brain. Symptoms of all brain tumors may produce that vary makes smaller or less in the amount on the part of the brain included. The condition of most brain tumors is familiar. This article will give an overview of brain tumors in population, summarized estimates of brain tumor symptoms, risk factors of brain tumors, brain tumors in children and young people.

Keywords: Factors, Brain tumor, Cell, Headache.

Abbreviation: WHO: World Health Organization, USA: United States of America, ICP: Influence of intracranial Pressure, CNS: Central Nervous System, CBT: Computer-based training.

INTRODUCTION

A brain tumor develops when unusual cells form in the brain. There are two class brain malignancies: cancerous (malignant) and benign (non- cancerous) tumors. cancerous tumors can be detached into primary tumors, which start inside the brain, and secondary tumors, which have spread from away, known as brain metastasis tumors[1]. Several assemblies are categorized neoplasms by primary and secondary brain tumors. There were 23,800 new cases of primary brain neoplasms (neoplasms are abnormal growth of tissues that can occur any part of the body). In the USA in 2017, it is estimated that one-fourth of patients with cancer will progress intracranial metastasis [2]. Secondary brain tumors are around four-times more communal than prime brain cancers [1], [3]. primary brain tumors happen in about 250,000 people a year worldwide, making up less than 2% of cancers [4]. About 1.2 million nervous system cancer is estimated in 2015 [5] 229.00 death occur due to cancer in 2015 [5]. In 2007, An American cancer civilization assessed that primary brain tumors were the cause of death in nearly 12,740 people, and 20,500 novel cases were diagnosed. According to the World Health Organization (WHO) classification, there are main classes of gliomas: astrocytomas, oligodendromas, and mixed oligoastrocytomas, which can characteristically be illustrious by their histological property [6].

Glioblastoma is the most malignant and most common glioma, accounting for 45-50% of gliomas [7].in 2008 there were calculated 1 52,236 new primary brain tumor (PBT) identification in the united states, with 22,000 being malignant (rate, 6/100,000 for a malignant tumor and 9/100,000 for being tumors). The general rate is slightly greater in females than males (17.2 vs 15.8/100,000) [8].

SYMPTOMS

There are following symptoms of the brain tumors

- ❖ problems Headaches as an influence of intracranial pressure (ICP) can be an early sign of brain cancer [9].In a sequence of 206 patients with brain tumors, 48% accessible with headache [10], [11]. Determinant conditions for presenting with headache were female gender, juvenile age and a history of headache .in particular of brain

tumor patients with long-established primary headache disorderliness history, 64% handover with headache [12]. while several articles statement on particulars of headache as a sign of brain tumors [10], [13].

- ❖ Two most normally happen indication in primary and solid brain tumor patients are fatigue and sleep-wake disruption [6], [12], [14], [15]. sleep-wake disorders are well-defined as a perceived or actual change in sleep that consequences in reduced daytime functioning, unlike other cancers. Sleep-wake disturbances are usual and are roughly calculate in 10%-20% of general population(paper=9)r17,18)[6], [16], [17].
- ❖ Symptoms of brain tumors occur under five-year preschool child are involve as Eye movement disorder, Co-ordination/walking problems, Behavior change, Seizures, Abnormal head position such as the wry neck, head, head tilt or stiff neck.
- ❖ 5-1-year kids head tumors take place symptoms to persevere/regular headache, persistent /recurrent vomiting, balance/co-ordination /walking problems, abnormal eye actions unclear or double vision, behavior alteration, seizures.
- ❖ In young people, brain tumors symptoms are repeated headache, persistent/recurrent vomiting, walking, Abnormal eye movements, blurred or double vision, behavior change, seizures, slow growth problems [18]–[20].
- ❖ Melancholy symptoms among brain tumor patients have been reported by a distinct investigative clinical interview with alternate criteria and doorstep [21], [22]. Expected range of depression is 2.8% to 95% [6], [21].

RISK FACTORS

Hereditary conditions such as Von Hippel-Lindau diseases, tuberous sclerosis, numerous endocrine neoplasias, and neurofibromatosis kind second transmit a high chance for the evolution of the brain tumors [1], [23], [24]. People with celiac disease have a slightly increased risk of developing brain tumors [25]. Environmental possibility factors for brain tumors inquire into epidemiological studies are ionizing radiation, mobile phone, Alcohol, chemical substances like hair dyes and pesticide, etc., “low-frequency” electromagnetic fields, Allergies[26]

BRAIN TUMORS OF CHILD

Brain tumors are the second most common cancer in children, comprising 15-25% of all pediatric malignancies, and they are the most common solid tumors. In the UK, around 450 childhood CNS tumors, which are approximately totally primary, investigated year influence children up to 15 years of age. They are the most usual solid tumor of childhood, including for 27% of all malignancies. participation of the brain is a risk for 55% of childhood malignancy, with 30% considered for by severe leukemias [17] [19]. Grad 1 gliomas are rare and mostly limited to childhood.

Diffuse astrocytomas (WHO grade II) description for 5% of all tumors in children aged 0- 14 years, with a U.S. occurrence rate of “28/100,000” [27]. High dose radiation to the head and neck for the action of cancer or other condition is to identify the CBT risk factor [2]. A Danish study surveyed CBT associated with neonatal diagnostic

X-ray submission and remarked a 2-fold positive non-significant suggestion [28]. Alcohol exposure in utero is a known toxin to progress CNS. But, in concurrence with past studies [29], a current large case-control study did not support maternal consumption as a risk factor for contribution CBTs[29] [29]. At the end of 2013, it was approximated that there were extra than 420000 childhood cancer survivors in the united state. childhood brain tumors stayers, henceforth mention to as survivors, representing 15.6% of all childhood cancer survivors [21].

BRAIN TUMORS IN YOUNG

Malignant tumors of the brain are a rare occurrence accounting for approximately 2% of all cancer in adults. Grad 2 gliomas (with subtypes astrocytoma and oligodendrogliomas) happen in all ages with a peak in young adults [6]. 13% of brain tumors all primary brain cancer present in persons younger than 20 years. [8], [30].In Finland, the antidepressant use in age groups 16 to 20 years has nearly tripled, to 6% and 3% from 1997 to 2007 [31]. In alternative Finnish study on the occurrence of depressive and anxiety signs in young adults, it was discovered that depressive disorders were the maximum usual mental disorder (6.5%), and they were more common in women (8.3% vs.4.6%)[32]. more exactly, symptoms influence poor movement, poor physical functioning, reduced bone mineral density, hearing and vision issues, and fitness levels [33].

CONCLUSION

The occurrence of new brain tumors 6.4 per 100,000 peoples per year with total five-year survival rate of 33.4%.there is an approximate 0.6% era risk of being analyzed with brain or other nervous system cancer[34].in childhood a different profile of tumor types is present and survival has improved over recent years and is higher than in adults.

REFERENCES

- [1] R. T. Merrell, "Brain Tumors," *YMDA*, vol. 58, no. 12, pp. 678–689, 2013.
- [2] B. D. Fox, V. J. Cheung, A. J. Patel, D. Suki, and G. Rao, "Epidemiology of Metastatic Brain Tu m or s," *Neurosurg. Clin. NA*, vol. 22, no. 1, pp. 1–6, 2011.
- [3] B. Connectivity, "Functional connectivity in adult brain tumor patients : A systematic review," no. 404, pp. 1–60, 2018.
- [4] S. Mcguire, "World Cancer Report 2014 . Geneva , Switzerland : World Health Organization , International Agency for Research on Cancer , WHO Press , 2015," pp. 418–419, 2015.
- [5] D. Collaborators, "Global , regional , and national life expectancy , all-cause mortality , and cause-specifi c mortality for 249 causes of death , 1980 – 2015 : a systematic analysis for the Global Burden of Disease Study 2015," pp. 1980–2015, 2016.
- [6] N. A. Access et al., "Neuro-Oncology," no. May, pp. 1–13, 2016.
- [7] K. Song, "Advance Care Planning in Patients with Primary Malignant Brain Tumors : A Systematic Review," vol. 6, no. October, 2016.
- [8] I. Review, "Brain Tumor Rehabilitation," vol. 90, no. 5, pp. 50–62, 2011.
- [9] K. Kahn and A. Finkel, "It IS a Tumor- Current Review of Headache and Brain Tumor," 2014.
- [10] T. Kurth, J. E. Buring, and P. M. Rist, "Headache , migraine and risk of brain tumors in women : prospective cohort study," 2015.
- [11] L. Valentinis et al., "Headache attributed to intracranial tumours : a prospective cohort study," no. January 2007, 2009.

- [12] J. N. Rich et al., “Statement brain tumours,” *Nat. Rev. Clin. Oncol.*, vol. 16, no. August, 2019.
- [13] L. P. Taylor, “Headache Currents Mechanism of Brain Tumor Headache,” pp. 772–775, 2014.
- [14] M. Dodd and R. N. Faan, “Advancing the science of symptom management,” 2001.
- [15] L. Rybicki, “Symptom clusters : myth or reality ?*,” vol. 24, no. June, pp. 373–385, 2010.
- [16] E. S. Ford, A. G. Wheaton, T. J. Cunningham, W. H. Giles, D. P. Chapman, and J. B. Croft, “Trends in Outpatient Visits for Insomnia , Sleep Apnea , and Prescriptions for Sleep Medications among US Adults : Findings from the National Ambulatory Medical Care Survey 1999-2010,” 2010.
- [17] Y. Dauvilliers and A. Buguet, “Clinical research,” pp. 347–356, 2005.
- [18] D. Walker, W. Hamilton, F. M. Walter, and C. Watts, “Strategies to accelerate diagnosis of primary brain tumors at the primary – secondary care interface in children and adults,” vol. 2, no. 2013, pp. 447–462.
- [19] E. Ward, C. Desantis, A. Robbins, B. Kohler, and A. Jemal, “Childhood and Adolescent Cancer Statistics , 2014,” vol. 64, no. 2, pp. 83–103, 2014.
- [20] E. Steliarova-foucher et al., “Geographical patterns and time trends of cancer incidence and survival among children and adolescents in Europe since the 1970s (the ACCIS project) : an epidemiological study,” pp. 2097–2105, 2004.
- [21] J. Huang, C. Zeng, J. Xiao, D. Zhao, and H. Tang, “Association between depression and brain tumor : a systematic review and meta-analysis,” vol. 8, no. 55, pp. 94932–94943, 2017.
- [22] A. Pranckeviciene and A. Bunevicius, “CNS Oncology Depression screening in patients with brain tumors : a review,” vol. 4, pp. 71–78, 2015.
- [23] T. S. Hodgson, S. M. Nielsen, M. S. Lesniak, and R. V Lukas, “Neurological Management of Von Hippel-Lindau Disease,” vol. 21, no. 5, pp. 73–78, 2016.
- [24] L. Rogers et al., “Meningiomas: knowledge base, treatment outcomes, and uncertainties. A RANO review,” 2014.
- [25] C. S. Hourigan, “The molecular basis of coeliac disease,” pp. 53–59, 2006.
- [26] R. A. Kleinerman et al., “Self-reported Electrical Appliance Use and Risk of Adult Brain Tumors,” vol. 161, no. 2, pp. 136–146, 2005.
- [27] K. M. Kurian et al., “Neuro-Oncology Practice,” vol. 5, no. August 2017, pp. 10–17, 2018.
- [28] L. Mellekjær, H. Hasle, G. Gridley, C. Johansen, S. K. Kjær, and K. Frederiksen, “Risk of cancer in children with the diagnosis immaturity at birth,” pp. 231–237, 2006.
- [29] C. Infante-rivard and M. El-zein, “Parental alcohol consumption and childhood cancers: a review,” pp. 101–129, 2007.
- [30] “SEER Cancer Statistics Review 1975-2005 National Cancer Institute,” p. 9989, 2005.
- [31] A. Palese, M. Cecconi, R. Moreale, and M. Skrap, “Pre-operative Stress , Anxiety , Depression and Coping Strategies Adopted by Patients Experiencing Their First or Recurrent Brain Neoplasm : An Explorative Study,” vol. 425, pp. 416–425, 2012.
- [32] A. Kero, M. Koivisto, L. Madanat-harjuoja, and N. Malila, “Purchases of antidepressants after cancer at a young age in Finland.”
- [33] E. Nicklin et al., “Long - term issues and supportive care needs of adolescent and young adult childhood brain tumour survivors and their caregivers : A systematic review,” no. January, pp. 477–487, 2019.
- [34] A. Perkins, G. Liu, and S. Alabama, “Primary Brain Tumors in Adults: Diagnosis and Treatment,” 2016.